

Utilisation of corn (*Zea mays*) bran and corn fiber in the production of food components[†]

Devin J Rose, George E Inglett* and Sean X. Liu

Abstract

The milling of corn for the production of food constituents results in a number of low-value co-products. Two of the major co-products produced by this operation are corn bran and corn fiber, which currently have low commercial value. This review focuses on current and prospective research surrounding the utilization of corn fiber and corn bran in the production of potentially higher-value food components. Corn bran and corn fiber contain potentially useful components that may be harvested through physical, chemical or enzymatic means for the production of food ingredients or additives, including corn fiber oil, corn fiber gum, cellulosic fiber gels, xylo-oligosaccharides and ferulic acid. Components of corn bran and corn fiber may also be converted to food chemicals such as vanillin and xylitol. Commercialization of processes for the isolation or production of food products from corn bran or corn fiber has been met with numerous technical challenges, therefore further research that improves the production of these components from corn bran or corn fiber is needed.

Published 2010 by John Wiley & Sons, Ltd.

Keywords: maize; co-products; corn fiber gum; vanillin; ferulic acid; xylo-oligosaccharides

INTRODUCTION

Corn (*Zea mays*) is a major crop in the USA, with 329×10^6 t produced domestically in 2008.¹ In comparison, 68.0×10^6 t of wheat and 9.24×10^6 t of rice were produced in 2008.^{2,3} Of the total domestic corn use, 52% was devoted to animal feed, 37% went to ethanol production (fuel and beverage) and 11% was allocated for the production of food products.⁴

The processing of corn for food use requires one of two milling techniques: dry-milling or wet-milling. Corn dry-milling involves the traditional milling of clean, tempered grain (moisture content 200 g kg^{-1}) to separate its component parts: endosperm, germ and bran.^{5,6} The major purpose of dry-milling is to recover the endosperm fraction for use as corn grits, meals and flours, while the germ may be harvested for oil.⁶ Corn bran, however, currently has low value and is often used for animal feed alone or in combination with corn germ cake or meal (germ after oil has been pressed or extracted). Wet-milling of corn involves first steeping the grain in water and sulfur dioxide. During this process the moisture content increases to about 450 g kg^{-1} and the kernels are softened to facilitate separation of the components: starch, gluten, fiber and germ.⁷ The profitable products of wet-milling are the starch (endosperm) and oil (germ) from the corn kernel. Co-products of wet-milling include corn fiber, corn gluten and steeping solids, which are sometimes combined and sold as corn gluten feed.⁷ Both corn bran and corn fiber are mainly composed of the pericarp (bran); however, corn fiber also contains cell wall material from the endosperm, which is not contained in corn bran.⁸

Corn bran is produced in yields of about $60\text{--}70 \text{ g kg}^{-1}$, while corn fiber is produced in yields of about $80\text{--}110 \text{ g kg}^{-1}$ total corn kernel.^{9,10} In 2008, 25.6×10^6 t of corn was wet-milled, while 5.24×10^6 t was dry-milled.⁴ Thus about 2.43×10^6 t of corn fiber and 0.341×10^6 t of corn bran were produced.

Traditional uses for these co-products of corn processing largely include animal feed, which does not command a high price. Therefore research is constantly under way to expand the use of these products. Lucrative applications of these products would provide much needed economic relief for farmers by increasing revenue and would benefit manufacturers and consumers by decreasing fuel and food costs. Much of the research concerning the utilization of corn bran and corn fiber has centered on its conversion to fuel ethanol;^{11,12} however, food applications of these co-products may also provide added value. Therefore in this review the composition of corn bran and corn fiber will be discussed and then current and prospective research surrounding the utilization of corn fiber and corn bran in the production of food components will be addressed.

COMPOSITION OF CORN BRAN AND CORN FIBER

The chemical composition of corn bran and corn fiber is shown in Table 1. Clearly, the majority of both corn bran and corn fiber is dietary fiber, which is nearly completely insoluble. The insoluble dietary fiber in corn bran is composed of cellulose ($\sim 280 \text{ g kg}^{-1}$) and hemicellulose ($\sim 700 \text{ g kg}^{-1}$), with only a small amount of

* Correspondence to: George E Inglett, Functional Foods Research Unit, National Center for Agricultural Utilization Research, USDA, ARS, 1815 N University Street, Peoria, IL 61604, USA. E-mail: George.Inglett@ars.usda.gov

† Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the US Department of Agriculture.

Functional Foods Research Unit, National Center for Agricultural Utilization Research, USDA, ARS, 1815 N University Street, Peoria, IL 61604, USA

Table 1. Composition (g kg⁻¹) of corn bran and corn fiber

Constituent	Corn bran	Corn fiber
Protein ^a	50–115 ^{13–15}	100–130 ^{16,17}
Starch	40–112 ^{13,14}	150–200 ^{16,17}
Oil	13.2–19 ^{14,18}	17.2–36.8 ¹⁸
Ferulate phytosterol esters ^b	0.2 ¹⁸	0.61–1.28 ¹⁸
Ash	6–10 ^{14,15}	6–20 ^{16,17}
Total dietary fiber	732–860 ^{13,19,20}	526–735 ²¹
Soluble fiber	2–26 ^{19,20}	ND–3 ²¹
Insoluble fiber	706–863 ^{19,20}	526–732 ²¹
Arabinose ^c	128–178 ^{13,14}	113–117 ^{12,16}
Xylose	217–243 ^{13,14}	176–213 ^{12,16}
Mannose	3 ¹⁵	ND–8.4 ²¹
Galactose	44–51 ^{13–15}	35.9 ¹⁶
Glucose ^d	182–248 ^{13,15}	300–372 ^{12,16}
Uronic acids	39–42 ^{13–15}	30–40 ¹⁷
Lignin	7–10 ^{13,14}	78 ¹⁶
Total phenolics	55 ¹³	NR
Ferulic acid	28–31 ^{13–15}	1.02–18.5 ^{22,23}
Diferulic acid	6.8–32 ^{15,24}	NR
<i>p</i> -Coumaric acid	3–4 ^{13,14}	2 ²²

ND, not detected; NR, not reported.

^a Nitrogen × 6.25.

^b Indentation indicates that this component is a component of the above constituent but is still reported as a proportion of the entire corn bran or corn fiber product.

^c Neutral sugars and uronic acids reported in polysaccharide form.

^d non-starch glucose.

lignin (~10 g kg⁻¹). The hemicellulosic fraction is composed of a β -(1 → 4)-xylopyranosyl backbone and α -L-arabinofuranosyl residues as side units linked (1 → 2) or (1 → 3) to the main chain.¹⁴ Side units of D-glucuronic acid or oligosaccharide side chains containing galactose, xylose and arabinose also occur.^{25,26} Some of the arabinofuranosyl side units contain a ferulic acid moiety esterified to O-5.^{13,27,28} Because of its complex structure, this hemicellulosic fraction may be referred to as a heteroxylan or, because the structure is mainly composed of xylosyl and arabinosyl units, an arabinoxylan. Saulnier and Thibault²⁹ elegantly described the interactions between these heteroxylans and other cell wall components within corn pericarp: the heteroxylan fraction fills the spaces between the cellulosic microfibrils, which are then crosslinked with each other and other cell wall components via di- and triferulate bridges, solidifying the cell wall's structure and rendering the heteroxylan insoluble.

The remainder of corn bran and corn fiber comprises residual starch, lipid, protein, ash, phenolic compounds and other trace phytochemicals (Table 1). Subtle differences in the amounts of these components are apparent. Most notably, corn fiber contains more than twice as much ferulate phytosterol esters, which can be recovered in corn fiber oil,¹⁸ while corn bran contains two times more ferulic acid.²³ The importance of these compounds will be discussed further in later sections.

UTILISATION OF WHOLE CORN BRAN AND CORN FIBER

Research on the utilization of corn-milling co-products in food products began with the addition of corn bran to breads,³⁰ cakes³¹

and muffins³² with the aim of increasing the dietary fiber content of widely consumed foods. Unfortunately, this resulted in undesirable changes in product quality: bread containing corn bran at 200 g kg⁻¹ flour showed a 20% reduction in loaf volume,³⁰ layer cakes containing corn bran at 300 g kg⁻¹ flour showed decreases in most sensory scores, including texture, color and flavor,³¹ and corn bran at 250 g kg⁻¹ flour in muffins resulted in significant decreases in flavor, mouthfeel, texture and overall acceptability compared with muffins containing the same level of wheat bran.³²

In more recent experiments, Mendonca *et al.*³³ found that corn bran significantly decreased the radial expansion ratio, appearance and general acceptability of extruded snacks containing corn bran at 150–320 g kg⁻¹. Holguin-Acuna *et al.*³⁴ substituted corn bran for oat flour in an extruded breakfast cereal. They found that corn bran at 300 and 400 g kg⁻¹ gave the lowest breaking strength, and tested these two cereals in a sensory panel. Panelists preferred the cereal that contained corn bran at 300 g kg⁻¹. Unfortunately, the authors did not compare this sample with a control product containing no corn bran.

Artz *et al.*³⁵ hypothesized that the poor performance of corn bran in baked goods could be improved by extrusion.³⁶ Therefore they extruded corn bran using a twin-screw extruder³⁷ and then incorporated the extruded corn bran into cookies. Unfortunately, panelists were able to correctly identify the odd sample in triangle tests comparing control cookies with cookies containing extruded corn bran but were not able to correctly identify the odd sample in triangle tests comparing cookies containing extruded or unextruded corn bran,³⁵ indicating that extrusion did not change the baking properties of corn bran in this application.

In attempts to utilize other cereal brans, treatments such as careful control of particle size^{38,39} and enzyme treatment⁴⁰ have shown some promise; therefore these treatments with corn bran or corn fiber may also be beneficial, although perhaps not directly applicable.

De Kock *et al.*³⁸ and Ozturk *et al.*³⁹ demonstrated that bran particle size affects quality when added to bread and cookies. In bread, smaller particle sizes decreased loaf volume more than larger particle sizes.³⁸ In cookies, medium (212–425 μ m) and coarse (425–850 μ m) particle sizes gave better spread ratios, color and overall sensory scores compared with cookies made with finer particle sizes (<212 μ m).³⁹

Enzyme treatment may also affect the quality of baked goods with added cereal brans. The best example of this is the use of *endo*-(1 → 4)- β -xylanases (xylanases; EC 3.2.1.8) in bread making. During bread making, the presence of insoluble arabinoxylans increases water absorption^{41–43} and decreases loaf volume,⁴⁴ thus diminishing bread quality. Soluble arabinoxylans have a positive impact on bread quality owing to their contribution to dough viscosity, air entrapment and the improvement of loaf volume and texture.^{41,45–47} Xylanase addition to bread dough can convert a portion of the insoluble arabinoxylans to soluble oligosaccharides and improve loaf volume.^{48,49} Because corn bran and corn fiber contain substantial levels of arabinoxylans (Table 1), adding xylanase to bread fortified with corn bran or corn fiber may improve its loaf volume and sensory attributes. This may also extend to other products such as cookies.⁴⁰

ISOLATION AND UTILISATION OF COMPONENTS OF CORN BRAN AND CORN FIBER

Corn bran and corn fiber contain potentially valuable components, including oil, phytosterol esters, dietary fiber and phenolic

antioxidants. Therefore some researchers have explored the use of processing technologies to harvest these components. Research on the extraction, production and utilization of each of these components will be discussed separately.

Cellulosic fiber gel

Corn bran contains about 200 g cellulose kg⁻¹ (Table 1). Usually, cellulose exhibits undesirable characteristics in foods and has few desirable functional properties;⁵⁰ however, the native structure may be disrupted in a two-stage high-shear process to yield a cellulosic fiber gel.^{51,52} The first stage involves treatment with a base to completely disintegrate the cell wall structure of corn bran. The solids are then recovered and treated again with hydrogen peroxide to produce a colorless product. Cellulosic fiber gel from corn bran exhibits high hydration capacity, high viscosity and a gel-like structure.⁵³

Cellulosic fiber gel from corn bran has found commercial success. Marketed under the trade name Z-Trim, it is promoted as a fat mimetic or flour substitute and is used in baked goods, condiments, dairy foods and processed meats (<http://www.ztrim.com>). Furthermore, Z-Trim is finding applications in school lunches to help meet the strict nutritional guidelines outlined in the National School Lunch Program set forth by the Food and Nutrition Service of the US Department of Agriculture.⁵⁴

One example of Z-Trim's success was reported by Warner and Inglett.⁵⁵ Brownies prepared with Z-Trim containing 40% less fat than control brownies and 50% flour replacement were evaluated by trained panelists. The Z-Trim brownies were rated as not significantly different from the control brownies with respect to chocolate flavor, sweetness, bitter flavor, staleness and cereal flavor. In addition, the Z-Trim brownies rated significantly higher for moistness, density and cohesiveness, which are desirable characteristics in a fudge-like brownie.

Corn fiber gum

As mentioned, the dietary fiber fraction in both corn bran and corn fiber is composed largely of heteroxylan, which is insoluble owing to its di- and triferulate crosslinks.²⁸ A substantial portion of this heteroxylan may be removed by treatment of corn fiber or corn bran with alkali or alkaline hydrogen peroxide followed by acidification to pH 4.0–4.5 (to remove insoluble hemicellulose A) and ethanol precipitation.^{10,14} The product thus obtained has been termed corn fiber gum.²⁶

A myriad of treatment conditions have been used to solubilize corn fiber gum. Chanliaud *et al.*¹⁴ used response surface experimental design in two experiments studying the effects of alkali type and concentration, time, temperature and liquid/solid ratio on corn fiber gum yields. When using potassium hydroxide, maximum yields of about 870 g kg⁻¹ of available heteroxylan were obtained with alkali concentrations between 1.2 and 1.5 mol L⁻¹ at 100 °C. Using saturated calcium hydroxide, maximum yields similar to those with potassium hydroxide were obtained after 19 h at 97 °C.

Hespe¹⁷ found that potassium hydroxide and ammonium hydroxide produced dark extracts, probably due to colored compounds produced at the high temperatures used for extraction. This phenomenon was also shown with sodium hydroxide.⁵⁶ These dark-colored heteroxylan products would likely be undesirable owing to their influence on product color.¹⁰ Calcium hydroxide produced a corn fiber gum that was free of impurities, lighter in color and dissolved easily in water.¹⁷ Unfortunately, yields were

low (<380 g kg⁻¹ of available heteroxylan) and the cellulose-rich residue remaining after corn fiber gum extraction was hard and unusable. Higher yields with greater purity were obtained when corn fiber gum was extracted with potassium hydroxide and then the extract was treated with calcium hydroxide.

Another approach to producing a lighter-colored corn fiber gum has been to use hydrogen peroxide. This converts much of the colored lignin compounds to soluble organic acids that can be removed.⁵⁷ Delignification occurs optimally at pH 11.5;⁵⁷ therefore hydrogen peroxide can be added to the alkali during extraction¹⁰ or the solubilized corn fiber gum may be treated with hydrogen peroxide after extraction,⁵⁸ the latter being more efficient.

As stated previously, corn fiber includes cellular material from the whole grain, while corn bran is composed mainly of the pericarp and does not include the cell wall material from the endosperm. Thus differences in structure or physical properties may occur between corn fiber gums isolated from these two milling co-products, depending on whether heteroxylans from the endosperm have different characteristics than those in the pericarp. Yadav *et al.*^{59,60} isolated corn fiber gum from corn bran and corn fiber using the same extraction procedure, allowing for direct comparison. They boiled each starting material in dilute sodium hydroxide and calcium hydroxide. The soluble extract was then treated with hydrogen peroxide and partially acidified and the corn fiber gum was recovered by ethanol precipitation. They termed this isolate CFG-1. The residue left after dilute alkali extraction was further extracted with a more severe alkaline hydrogen peroxide treatment to obtain CFG-2.

Few differences in chemical composition and physical properties between these four fractions (CFG-1 and CFG-2 from corn bran and corn fiber) were evident.^{59,60} The molecular weights of the two CFG-1s were 290 000 and 334 000 Da when isolated from corn bran and corn fiber respectively, while the molecular weights of the two CFG-2s were 491 000 and 452 000 Da respectively. Corn fiber gums isolated from both sources showed low viscosities, with CFG-1 and CFG-2 from corn bran being slightly lower than CFG-1 and CFG-2 from corn fiber. The arabinose/xylose ratios of CFG-1 and CFG-2 isolated from corn bran were 0.55 and 0.60 and from corn fiber 0.70 and 0.67 respectively. This suggests that heteroxylans originating from the endosperm are slightly more branched than those from the pericarp owing to the higher arabinose/xylose ratios in heteroxylans from corn fiber (which contained endosperm heteroxylans) compared with corn bran (which did not contain endosperm heteroxylans). This was further substantiated when the heteroxylan fraction was isolated from coarse corn fiber (corn fiber without the endosperm fraction), which gave arabinose/xylose ratios of 0.60 and 0.64 for CFG-1 and CFG-2 respectively.⁵⁹ Therefore subtle differences in heteroxylan branching may exist between endosperm and pericarp heteroxylans, although these minor differences are certainly not as varied as for heteroxylans from other grains such as wheat.⁶¹

Perhaps the most substantial difference between the corn fiber gums isolated from these two co-products was that both CFG-1 and CFG-2 from corn fiber were significantly better emulsifiers than the same products isolated from corn bran. In fact, Yadav *et al.*⁵⁹ showed that these CFGs were better emulsifiers than gum arabic in orange oil emulsions. They attributed this to the higher protein content of the samples originating from corn fiber. This suggests that proteins may be more intimately associated with heteroxylans in the endosperm than in the pericarp.

Recently, Carvajal-Millan *et al.*²⁴ and Yadav *et al.*⁶² showed that some ferulic acid remains esterified to solubilized corn fiber gum

after alkali treatment. This is an important finding for two reasons. First, in the presence of hydrogen peroxide and peroxidase, corn fiber gum that contains ferulic acid can form gels via oxidative crosslinking, which can substantially affect its physical properties and potential applications.⁶³ Second, ferulic acid has received considerable attention owing to its antioxidant properties, and ferulic acid bound to soluble corn bran gum may be particularly important for delivering beneficial antioxidants to the colon for disease prevention. Indeed, while free phenolics are rapidly absorbed in the upper gastrointestinal tract,⁶⁴ bound phenolics can be released by microbial esterases in the lower gastrointestinal tract⁶⁵ and provide radical-scavenging activity in this region of the colon,⁶⁶ a region chronically under oxidative attack^{67,68} and prone to disease.⁶⁹

Corn fiber gum has not enjoyed the commercial success that has been afforded cellulosic fiber gel (discussed above). Because this product has several promising characteristics similar to those of exudate gums,⁷⁰ such as emulsification⁶⁰ and gel formation,⁶³ as described above, more research aimed at broadening the applications of this polysaccharide gum may eventually lead to commercialization as the market base widens. New applications could explore its use in expanded snacks, baked goods, beverages, specialty foods, edible coatings, supplements and the like.

Xylo-oligosaccharides

Xylo-oligosaccharides (XOS) are partially hydrolyzed, water-soluble xylan fragments that may be obtained from corn bran, corn fiber and other agricultural products by enzymatic or high-temperature treatment.^{71,72} Enzymatic treatment proceeds by utilizing microbial xylanases, while high-temperature processing employs hydronium ions and organic acids generated *in situ* at high processing temperatures (160–220 °C) to partially hydrolyze heteroxylan polymers and yield soluble hydrolysates.^{71,72}

Few studies exist on the use of corn bran or corn fiber as starting material for the enzymatic production of XOS. Research on the production of XOS from other cereal brans has revealed that 250–450 g kg⁻¹ of available heteroxylan may be released from the starting material with xylanase treatments, depending on processing conditions.^{73,74} However, previous researchers have shown that corn bran or corn fiber heteroxylan is a more difficult substrate for enzymatic degradation compared with others such as wheat bran.²³ Therefore new xylanases that are more specific for corn heteroxylan are needed.

One of the most important factors to consider in the enzymatic production of XOS from agricultural products is substrate selectivity of the xylanase. Direct enzymatic hydrolysis requires xylanases with high substrate selectivity toward insoluble xylan. Moers *et al.*⁷⁵ screened xylanases from *Bacillus subtilis*, *Aspergillus aculeatus*, *Aspergillus niger*, *Trichoderma viride* and *Trichoderma longibrachiatum* for their activities on soluble and insoluble xylan substrates. They found that the enzyme preparations from *B. subtilis* and *A. niger* were the most active on insoluble substrates, while those from *A. aculeatus* and *T. viride* were the most active on soluble xylan. Xylanases are also classified into a number of families based on their catalytic activity.⁷⁶ Xylanases from families 10 and 11 are active on heteroxylans such as those in corn bran.⁷⁶ For the production of XOS, family 11 may be more desirable owing to lower production of xylose,⁷⁷ although family 10 is more able to hydrolyze highly branched arabinoxylans.⁷⁸

Some researchers have shown that the enzymatic production of XOS can be increased by adding accessory enzymes. For instance, de Vries *et al.*⁷⁹ showed that arabinofuranosidases (EC

3.2.1.55) from *Aspergillus* fungi resulted in a 2.0–2.5-fold increase in the release of xylan. A feruloyl esterase (EC 3.1.1.73), which cleaves the ester linkages between ferulic acid and arabinosyl units on heteroxylans, may also increase enzymatic accessibility of the xylanase toward the heteroxylan fraction, thus releasing more XOS,^{80,81} although this has not been shown in all cases.⁸² While more XOS may be released with the addition of these enzymes, their structures and consequently their prebiotic and/or antioxidant properties may change substantially.

Autohydrolytic treatment has the advantage over enzymatic treatment in that it does not require the use of harsh chemicals or enzymes that can contribute substantially to cost.⁷² Unfortunately, autohydrolysate liquors contain a variety of contaminants that must be removed. These contaminants are commonly removed by activated charcoal, ultrafiltration, ion exchange, etc.⁸³

Two studies have reported the use of autohydrolysis to solubilize a portion of the heteroxylan fraction from corn bran or corn fiber.^{84,85} When corn fiber was used as a starting material, a maximum solubilization of 300 g kg⁻¹ of xylan occurred at 210 °C, but a substantial amount of monosaccharides was produced.⁸⁴ At 180 °C, xylan yield was lower (200 g kg⁻¹), but it contained a more desirable range of molecular weights. Using corn bran as a starting material, maximum solubilization of xylan occurred at 180 °C for 10 min or 200 °C for 2 min.⁸⁵ Under these conditions, about 500 g kg⁻¹ of arabinoxylan was released into solution in the form of oligo- and polysaccharides. These liquors were accompanied by a monosaccharide contamination of about 100 g kg⁻¹ of total neutral sugar residues in the starting material. These studies suggest that corn bran may be a better candidate for the production of XOS by autohydrolysis owing to higher yields.

Short-chain XOS from other agricultural products are potential prebiotics.⁸⁶ Prebiotics are indigestible oligosaccharides that selectively stimulate the growth of beneficial bacteria (e.g. bifidobacteria and lactobacilli) in the gastrointestinal tract of healthy persons.⁸⁷ In preliminary animal studies, XOS have been shown to stimulate the growth of bifidobacteria more than fructo-oligosaccharides, the most common and studied prebiotics.⁸⁸ In a study with aged men using a commercial XOS (Xylooligo 95P, Suntory Ltd, Osaka, Japan), concentrations of bifidobacteria were increased after 3 weeks on XOS.⁸⁹ XOS from corn bran or corn fiber exhibit similar structures to these prebiotic XOS, and this may also confer prebiotic properties.

In addition to prebiotic activity, XOS may contain antioxidant activity. Depending on production parameters, XOS produced from cereal brans may retain esterified ferulic acid moieties on the solubilized oligosaccharides. These XOS exhibit high antioxidant activity *in vitro*^{90–92} and, despite the antimicrobial effects of ferulic acid,⁹³ do not appear to inhibit the growth of probiotic bacteria in pure culture.⁹⁴ In the large bowel of rats, microbial esterases are able to release the ferulic acid moiety, where it acts as a local antioxidant or may be absorbed and transported to other tissues.⁹⁵ This has potential implications for the prevention of inflammatory diseases and cancer.⁶⁶ Notably, corn bran contains more ferulic acid than many common cereal brans, fruits and vegetables.⁹⁶ Thus XOS with bound ferulic acid from corn bran have the potential to exhibit higher antioxidant activity than those isolated from other agricultural products.

Corn fiber oil

Most corn oil for human consumption comes from the germ;⁹⁷ however, in recent years, corn fiber oil has received attention owing to its ability to lower cholesterol in animal studies.^{98–100}

This is most likely a result of high levels of ferulate phytosterol esters,¹⁰¹ the most predominant being sitostanyl ferulate.¹⁰² Corn fiber contains three to six times more of these components than corn bran (Table 1). Unfortunately, the amounts are still quite low compared with the levels of similar type compounds in other bran-derived oils such as rice bran oil.¹⁰² Therefore, before this product can obtain commercial application, ways to increase the oil yield and phytosterol content of the oil must be examined.

Singh *et al.*¹⁰³ explored the effects of adding different sulfates or acids to the steep water during the wet-milling of two hybrids of corn. None of the sulfates increased oil yield, but ammonium sulfate increased ferulate phytosterol ester yield. Of the acids tested, acetic and hydrochloric acids appeared to be the most effective at increasing oil yield and phytosterol content of the oil.

Acids or enzymes added to the corn fiber itself may affect oil yield and composition. Singh *et al.*¹⁰⁴ treated corn fiber that had been conventionally wet-milled with sulfuric acid or a mixture of cellulase, amylase, xylanase and β -glucosidase. They found that each of these treatments increased the oil and phytosterol content of the corn fiber by up to sevenfold by hydrolyzing and removing a portion of the non-oil components (e.g. starch, cellulose, hemicellulose).

Non-chemical treatments to increase oil yield of corn fiber have been investigated. Wu and Norton¹⁰⁵ used fine grinding and air classification to fractionate corn fiber. They found that the finest particles ($<30\ \mu\text{m}$) were highest in ferulate phytosterol esters. Moreau *et al.*¹⁰⁶ tested the effects of heat treatments in conventional, vacuum and microwave ovens. They found that none of the treatments substantially impacted oil yield but that the conventional and vacuum oven treatments slightly decreased phytosterols and substantially increased γ -tocopherol. They concluded that the decreases in phytosterols were practically insignificant and that the treatments were useful for increasing γ -tocopherol (an antioxidant) in corn fiber oil.

Ferulic acid

Ferulic acid is a phenolic antioxidant.¹⁰⁷ In comparison with other naturally occurring antioxidants, including gallic acid, caffeic acid, malvidin, delphinidin, catechin, epicatechin, rutin and quercetin, ferulic acid has been shown to be the most efficient in inhibiting lipid and protein oxidation in a lecithin/liposome oxidation system.¹⁰⁸ Ferulic acid also possesses antimicrobial activity against spoilage and pathogenic micro-organisms.^{109,110} Because of its antioxidant and antimicrobial properties, ferulic acid has great potential for use in the food industry as a preservative.¹¹¹

Ferulic acid may also be used in the production of vanillin, in edible films, as a crosslinking agent and as a supplement.¹¹¹ The use of ferulic acid as a precursor for natural vanillin production will be discussed in a later section. In edible films, ferulic acid has been shown to increase tensile strength and per cent elongation at break and decrease water vapor permeability and gas permeability in edible films created from soy protein isolate.¹¹² Ferulic acid addition to milk has been shown to enhance heat stability, possibly by crosslinking nucleophilic amino acid residues on κ -casein, thus preventing dissociation.¹¹³ As a supplement or drug, ferulic acid may theoretically be useful in the treatment of Alzheimer's disease, diabetes, some cancers, hypertension, atherosclerosis and inflammatory diseases.⁹⁶

Commercially, ferulic acid can be synthesized by a condensation reaction of vanillin with malonic acid catalyzed by piperidine.¹¹⁴ However, the cost of this procedure, environmental concerns and the desire for naturally derived food additives make isolation of

ferulic acid from natural sources desirable. Indeed, ferulic acid has been produced commercially in Japan by saponification of the product of rice bran oil refining¹⁰⁷. Interestingly, rice bran oil contains $10\text{--}20\ \text{g kg}^{-1}$ ferulic acid,^{107,111} whereas corn bran contains more than $30\ \text{g kg}^{-1}$ ferulic acid (Table 1) and is the best source of ferulic acid among common cereals, fruits, vegetables and other agricultural products.⁹⁶ Unfortunately, ferulic acid in corn bran is largely bound to cell wall components,^{27,28} thus creating technological challenges for its isolation from corn bran.

Much of the research surrounding ferulic acid utilization from corn bran has focused on its release from its bound state. In a process similar to the production of corn fiber gum (discussed above), corn bran may be treated with alkali to cleave the ferulate crosslinks,¹³ thus releasing ferulic acid into solution. In this case, ferulic acid and corn fiber gum are co-solubilized and can be separated by precipitating the corn fiber gum with ethanol.¹³ Saulnier *et al.*¹³ found a maximum release of ferulic acid from corn bran with $0.5\ \text{mol L}^{-1}$ sodium hydroxide at $30\ ^\circ\text{C}$ for 2 h, while only 30% of the corn bran gum fraction was co-solubilized. Thus, if corn bran gum is also desired, a harsher alkali treatment is required to release the heteroxylan fraction in higher yields.¹⁴ Tilay *et al.*¹¹⁵ used response surface methodology to maximize the release of ferulic acid from corn bran. They found that $4\ \text{mol L}^{-1}$ sodium hydroxide at $21.6\ ^\circ\text{C}$ for 24 h led to a 1.3-fold increase in the level of ferulic acid recovered compared with the unoptimized procedure, which was $2\ \text{mol L}^{-1}$ sodium hydroxide at room temperature for 24 h. The authors reported ferulic acid at $191.1\ \text{g kg}^{-1}$ corn bran under the optimized conditions. These levels are much lower than typically reported for corn bran, but comparable to corn fiber (Table 1). Therefore these authors may unknowingly have used corn fiber instead of corn bran, or the corn bran may have been old and undergone significant storage-related oxidation. The fact that the optimized extraction conditions are so different from those reported previously by Saulnier *et al.*¹³ suggests the former explanation.

Ferulic acid can also be liberated from corn bran by microbial enzymes, namely feruloyl esterases.^{116–118} Feruloyl esterases catalyze the cleavage of the ester linkage between ferulic acid and the arabinosyl moiety on arabinoxylans. These extracellular enzymes are produced from a number of bacteria and fungi when grown on substrates containing bound ferulic acid.^{116,118,119} Feruloyl esterases have been classified into types A, B, C and D based on their specificity for aromatic substrates and sequence similarities with other enzyme classes.¹¹⁶ Types A and D are able to cleave ferulic acid dimers such as those that form the crosslinks in corn pericarp cell walls, while all classes are able to release ferulic acid. Type A shares sequence similarity with lipases, while type B is similar to carboxylic esterase family 1. Types C and D show sequence similarities with chlorogenate esterase and xylanase respectively.

Microbial feruloyl esterases alone are not sufficient to release significant amounts of ferulic acid from corn bran;¹²⁰ other treatments and cell wall-degrading enzymes must be used in concert with feruloyl esterases if sufficient yields are to be realized. For example, Bonnin *et al.*¹⁵ were only able to release $30\ \text{g kg}^{-1}$ of insoluble, esterified ferulic acid from corn bran when incubated with an enzyme preparation from *A. niger* I-1472. However, when the corn bran was treated by autoclaving at $160\ ^\circ\text{C}$ for 1 h, the enzymes were able to release $903\ \text{g kg}^{-1}$ of esterified ferulic acid from the fraction of corn bran that had been solubilized by autoclave treatment. Using Novozym 342, Saulnier *et al.*¹²¹ solubilized only $300\ \text{g kg}^{-1}$ of insoluble ferulic acid from corn

bran, and about one-third of the ferulic acid remained esterified to arabinose. Upon pre-treating the corn bran by autoclaving at 160 °C for 1 h or flash-explosion at 190 °C for 1 min, 800 g kg⁻¹ of insoluble ferulic acid was solubilized (free + esterified) and Novozyme 342 was then able to release about two-thirds of that ferulic acid as free ferulic acid. When feruloyl esterases are used in combination with other enzymes, the increased release of ferulic acid is the result of the production of short-chain feruloylated XOS, which are soluble and more accessible to feruloyl esterases.¹²²

The above studies used a mixture of cell wall-degrading enzymes in addition to pre-treatments of the corn bran in order to facilitate enzymatic attack.^{15,121} This is because, in general, corn bran has proven a more difficult substrate for enzymatic degradation compared with others such as wheat bran.²³ Using a crude enzyme preparation from *Neosartorya spinosa* NRRL 185, however, Shin *et al.*²³ were able to release 988 g kg⁻¹ of insoluble, esterified ferulic acid as free ferulic acid from corn bran without pre-treatment.

BIOCONVERSION OF CORN BRAN AND CORN FIBER COMPONENTS

Most of the research on bioconversion of corn bran and corn fiber has focused on the production of fuel ethanol;^{11,12} however, components of these co-products may be converted to other chemicals that would be important to the food industry. The use of corn bran or corn fiber as a starting material for food additive/chemical production has numerous potential advantages such as reduced environmental impact, 'natural' label and increased revenue for producers of corn products.

Xylitol

Xylitol is a five-carbon polyalcohol that occurs widely in nature.¹²³ In the food industry, xylitol is used as a low-calorie, non-cariogenic sweetener with bulking properties similar to those of sucrose.¹²⁴ Xylitol is not used by plaque-causing bacteria in the mouth and thus does not promote dental caries; it is also useful for diabetic foods owing to its slow absorption and low impact on glycemic index.¹²⁴

Commercially, the majority of xylitol is produced by chemical hydrogenation of xylose, usually obtained from wood hydrolysates. Corn bran and corn fiber offer promising starting materials for xylose harvest owing to their high xylose content (~200 g kg⁻¹, Table 1), which is comparable to that of hardwoods.¹²⁵

Xylose from acid hydrolysates must be purified prior to chemical hydrogenation, because the presence of other materials impedes the conversion and crystallization of xylose.¹²⁴ This costly and technically challenging step, as well as environmental concerns related to chemical hydrogenation, has sparked interest in the microbial conversion of xylose to xylitol in a process similar to the production of ethanol. In this case the starting material is hydrolyzed using dilute acid or enzyme treatment and then the resulting syrup is fermented with a yeast such as *Candida tropicalis*.¹²⁶ As with the conversion of pentose sugars to ethanol, yields of xylitol are generally low because of the formation of inhibitors such as furfural and hydroxymethylfurfural during hydrolysis.¹²⁶ To overcome the effects of inhibitors, Buhner and Agblevor¹²⁷ treated corn fiber hydrolysates with activated charcoal to remove these inhibitors. They found that activated charcoal was effective at removing inhibitors and increasing xylitol yield. Improved processes and yeasts for the conversion of xylose to xylitol would be important.

Vanillin

Vanillin is the major flavor component in properly cured vanilla (*Vanilla planifolia* Andrews) pods and occurs at 10–30 g kg⁻¹ whole bean.¹²⁸ Vanillin also occurs at lower concentrations in a wide variety of other agricultural products, including ponderosa pine (*Pinus ponderosa*), tobacco and citrus fruits;^{129,130} even corn bran itself contains a small amount of vanillin (55 g kg⁻¹).¹³¹

In the food, pharmaceutical and cosmetic industries, vanillin is an important flavor and aroma compound.¹³² In these industries the demand for vanillin far outweighs its production from vanilla beans; therefore synthetic (or artificial) vanillin is often used.¹²⁸ Between the 1930s and 1980s in the USA, most of the synthetic vanillin was produced from spent sulfite liquors; however, owing to environmental concerns, this process has been abolished.¹²⁹ Today, synthetic vanillin is produced from guaiacol using petroleum as the carbon source.^{128,129} Owing to the desire for natural vanillin production with less environmental impact, researchers have turned to ferulic acid as a possible starting material, which can be converted to vanillin by microbiological means.^{133–135} Hence, owing to its high ferulic acid content, corn bran may play a critical role in supplying ferulic acid for the biosynthesis of vanillin.

As mentioned, the ferulic acid in corn bran is covalently linked to arabinosyl moieties as a glycosidic conjugate^{27,28} and thus must be liberated from its bound state. Alkaline hydrolysis can accomplish this task;^{13,115} however, ferulic acid obtained in this manner cannot be considered natural.¹³⁶ For a natural claim, ferulic acid and other potential food additives must be obtained by physical, enzymatic or microbiological means.¹³⁷

Lesage-Meessen *et al.*¹³⁷ used a three-step process to convert ferulic acid from corn bran to vanillin. The first step involved autoclaving corn bran at 140 °C for 3 h and recovering the solubilized feruloylated oligosaccharides, which represented 300 g kg⁻¹ of total ferulic acid originally present in the corn bran. Enzymes harvested from *A. niger*, which had been grown on sugar beet pulp, were then added to a culture of *Pycnoporus cinnabarinus* containing the soluble feruloylated oligosaccharides obtained from autoclaved corn bran. The *A. niger* enzymes released the ferulic acid from its carbohydrate moiety and converted the free ferulic acid to vanillic acid; *P. cinnabarinus* then converted the vanillic acid to vanillin with a molar yield of 22%. When *A. niger* itself, rather than its enzymes, was used to release ferulic acid and convert it to vanillic acid, followed by conversion with *P. cinnabarinus*, the molar yield increased to 43% when cellobiose and XAD-2 resin were also added to the culture medium.

As discussed previously, enzyme extracts from *N. spinosa* NRRL 185 were able to release 988 g kg⁻¹ of total ferulic acid as free ferulic acid from corn bran without pre-treatment.²³ The ferulic acid thus obtained was then added to a culture of *Streptomyces setonii* ATCC 391 161. After 12 h, 43% of the ferulic acid had been converted to vanillin; however, substantial levels of vanillic acid and other contaminants were also produced.

Buranov and Mazza¹³¹ recently reported a possible non-microbiological method of converting ferulic acid to vanillin that would still be considered natural. This involves the direct conversion of bound ferulic acid to vanillin using pressurized low-polarity water extraction (also known as subcritical water extraction). In this process, water at high temperature was kept under sufficient pressure to maintain it in its liquid state while passing through a column containing the sample. Buranov and Mazza¹³¹ hypothesized that, under these conditions, some of the ferulic acid was liberated from its bound state by cleavage

of its aliphatic double bond rather than the ester linkage, thus producing vanillin. The highest vanillin production (1.68 g kg⁻¹ corn bran) was obtained using water at 220 °C, 5.2 MPa and a flow rate of 5 mL min⁻¹ for 40 min. This represents a poor yield, as the corn bran starting material contained 25.1 g ferulic acid kg⁻¹. The authors suggest that yields may be improved by optimizing processing conditions and/or adding 'green' additives.

CONCLUSIONS

Wet-milling and dry-milling of corn result in a variety of co-products that are currently sold as animal feeds and other low-value commodities.^{6,7} Utilization of these co-products for food applications may increase the value of these mill streams, reduce agricultural waste and decrease the cost of major corn products. Two of these major co-products are corn bran and corn fiber.

Previous research has been devoted to direct addition of these mill streams to food,^{30–34} fractionation of each co-product to recover potentially useful components^{8,10,14,17,24,52,58–60,84,85,101,103,104,106} or conversion of these components to other commercially important chemicals.^{23,127,131,137} Corn fiber oil, with its high levels of ferulate phyto-sterol esters, may help lower cholesterol,^{98–100} while phenolic acids, richly contained in corn bran, may be used as preservatives, as crosslinking agents, as supplements or as precursors to natural vanillin production.¹¹¹ The dietary fiber fraction of each co-product can be fractionated to yield corn fiber gum with low viscosity and excellent emulsifying properties^{59,60} or cellulosic fiber gels with properties similar to shortening in baked goods.^{52,55} The heteroxylan fraction can be partially hydrolyzed to yield potential prebiotics⁸⁶ or completely hydrolyzed and converted to food ingredients.¹²⁷

With the exception of cellulosic fiber gel (Z-Trim) produced from corn bran,⁵² the commercial implementation of these applications has not been realized owing to the numerous challenges involved in working with these co-products. Important areas of future research may include the following: (1) improved techniques beyond extrusion for increasing functionality of these whole co-products in foods; (2) ways to increase corn fiber oil yield and maximize the phytosterol content; (3) improved methods for the solubilization or partial hydrolysis of the heteroxylans in corn bran or corn fiber to yield corn fiber gum or XOS; (4) more cost-effective ways of liberating and purifying ferulic acid and other phenolic acids; (5) improved methods for the complete hydrolysis of heteroxylans for conversion to xylitol or other food chemicals; (6) identification of more productive strains of micro-organisms that are capable of converting components of corn bran to potentially useful food products.

REFERENCES

- ERS, Table 4 – Corn: Supply and disappearance. [Online]. Available: <http://www.ers.usda.gov/Data/FeedGrains/Yearbook/FGYearbookTable04.pdf> [18 February 2010].
- ERS, Table 4 – World and U.S. wheat production, exports and ending stocks. [Online]. Available: <http://www.ers.usda.gov/Data/Wheat/Yearbook/WheatYearbookTable04.pdf> [18 February 2010].
- ERS, Table 1 – Rice supply and use. [Online]. Available: <http://www.ers.usda.gov/Briefing/Rice/data.htm> [21 May 2009].
- ERS, Table 31 – Corn: Food, seed, and industrial use. [Online]. Available: <http://www.ers.usda.gov/Data/FeedGrains/Yearbook/FGYearbookTable31.pdf> [18 February 2010].
- Burge RM and Duensing WJ, Processing and dietary fiber ingredient applications of corn bran. *Cereal Foods World* **34**:535, 537–538 (1989).
- Duensing WJ, Roskens AB and Alexander RJ, Corn dry milling: processes, products, and applications, in *Corn: Chemistry and Technology*, ed. by White PL and Johnson LA. American Association of Cereal Chemists, St Paul, MN, pp. 407–447 (2003).
- Johnson LA and May JB, Wet milling: the basis for corn biorefineries, in *Corn: Chemistry and Technology*, ed. by White PL and Johnson LA. American Association of Cereal Chemists, St Paul, MN, pp. 449–494 (2003).
- Singh V, Doner LW, Johnston DB, Hicks KB and Eckhoff SR, Comparison of coarse and fine corn fiber gum yields and sugar profiles. *Cereal Chem* **77**:560–561 (2000).
- Watson SA, Description, development, structure, and composition of the corn kernel, in *Corn: Chemistry and Technology*, ed. by White PL and Johnson LA. American Association of Cereal Chemists, St Paul, MN, pp. 69–106 (2003).
- Doner LW and Hicks KB, Isolation of hemicellulose from corn fiber by alkaline hydrogen peroxide extraction. *Cereal Chem* **74**:176–181 (1997).
- Dhugga KS, Maize biomass yield and composition for biofuels. *Crop Sci* **47**:2211–2227 (2007).
- Gaspar M, Kalman G and Reczey K, Corn fiber as a raw material for hemicellulose and ethanol production. *Proc Biochem* **42**:1135–1139 (2007).
- Saulnier L, Marot C, Chanliaud E and Thibault JF, Cell-wall polysaccharide interactions in maize bran. *Carbohydr Polym* **26**:279–287 (1995).
- Chanliaud E, Saulnier L and Thibault JF, Alkaline extraction and characterisation of heteroxylans from maize bran. *J Cereal Sci* **21**:195–203 (1995).
- Bonnin E, Saulnier L, Brunel M, Marota C, Lesage-Meessen L, Asther M, *et al*, Release of ferulic acid from agroindustrial by-products by the cell wall-degrading enzymes produced by *Aspergillus niger* I-1472. *Enzym Microb Technol* **31**:1000–1005 (2002).
- Grohmann K and Bothast RJ, Saccharification of corn fibre by combined treatment with dilute sulphuric acid and enzymes. *Process Biochem* **32**:405–415 (1997).
- Hespeil RB, Extraction and characterization of hemicellulose from the corn fiber produced by corn wet-milling processes. *J Agric Food Chem* **46**:2615–2619 (1998).
- Moreau RA, Powell MJ and Hicks KB, Extraction and quantitative analysis of oil from commercial corn fiber. *J Agric Food Chem* **44**:2149–2154 (1996).
- Kahlon TS and Chow FI, *In vitro* binding of bile acids by rice bran, oat bran, wheat bran, and corn bran. *Cereal Chem* **77**:518–521 (2000).
- Dikeman CL, Murphy MR and Fahey GC, Dietary fibers affect viscosity of solutions and simulated human gastric and small intestinal digesta. *J Nutr* **136**:913–919 (2006).
- Guevara MA, Bauer LL, Abbas CA, Beery KE, Holzgraefe DP, Cecava MJ, *et al*, Chemical composition, *in vitro* fermentation characteristics, and *in vivo* digestibility responses by dogs to select corn fibers. *J Agric Food Chem* **56**:1619–1626 (2008).
- Akin DE and Rigsby LL, Corn fiber: structure, composition, and response to enzymes for fermentable sugars and coproducts. *Appl Biochem Biotechnol* **144**:59–68 (2008).
- Shin HD, McClendon S, Le T, Taylor F and Chen RR, A complete enzymatic recovery of ferulic acid from corn residues with extracellular enzymes from *Neosartorya spinosa* NRRL 185. *Biotechnol Bioeng* **95**:1108–1115 (2006).
- Carvajal-Millan E, Rascon-Chu A, Marquez-Escalante JA, Micard V, Ponce de Leon N and Gardea A, Maize bran gum: extraction, characterization and functional properties. *Carbohydr Polym* **69**:280–285 (2007).
- Montgomery R and Smith F, Structure of corn hull hemicellulose. III. Identification of the methylated aldobiuronic acid obtained from methyl corn hull hemicellulose. *J Am Chem Soc* **79**:695–697 (1957).
- Whistler RL and Corbett WM, Oligosaccharides from partial acid hydrolysis of corn fiber hemicellulose. *J Am Chem Soc* **77**:6328–6330 (1955).
- Saulnier L, Vigouroux J and Thibault JF, Isolation and partial characterization of feruloylated oligosaccharides from maize bran. *Carbohydr Res* **272**:241–253 (1995).

- 28 Saulnier L, Crepeau MJ, Lahaye M, Thibault JF, Garcia-Conesa MT and Kroon PA, Isolation and structural determination of two 5,5'-diferulate oligosaccharides indicate that maize heteroxylans are covalently cross-linked by oxidatively-coupled ferulates. *Carbohydr Res* **320**:82–92 (1999).
- 29 Saulnier L and Thibault JF, Ferulic acid and diferulic acids as components of sugar-beet pectins and maize bran heteroxylans. *J Agric Food Chem* **79**:396–402 (1999).
- 30 Sosulski FW and Wu KK, High-fiber breads containing field pea hulls, wheat, corn, and wild oat brans. *Cereal Chem* **65**:186–191 (1988).
- 31 Schafter MAM and Zabik ME, Dietary fiber sources for baked products: comparison of wheat brans and other cereal brans in layer cakes. *J Food Sci* **43**:375–379 (1978).
- 32 Polizzoto LM, Tinsley AM, Weber CW and Berry JW, Dietary fiber in muffins. *J Food Sci* **48**:111–113, 118 (1983).
- 33 Mendonca S, Grossmann MVE and Verhe R, Corn bran as a fibre source in expanded snacks. *Lebensm Wiss Technol* **33**:2–8 (2000).
- 34 Holguin-Acuna AL, Carvajal-Millan E, Santana-Rodriguez V, Rascon-Chu A, Marquez-Escalante JA, Ponce de Leon-Renova NE, et al, Maize bran/oat flour extruded breakfast cereal: a novel source of complex polysaccharides and an antioxidant. *Food Chem* **111**:654–657 (2008).
- 35 Artz WE, Warren CC, Mohring AE and Villota R, Incorporation of corn fiber into sugar snap cookies. *Cereal Chem* **67**:303–305 (1990).
- 36 Fulger L and Bradbury G, Modification of bran by extrusion. US Patent 4500558 (1985).
- 37 Artz WE, Warren C and Villota R, Twin screw extrusion modification of a corn fiber and corn starch extruded blend. *J Food Sci* **55**:746–754 (1990).
- 38 De Kock S, Taylor J and Taylor JRN, Effect of heat treatment and particle size of different brans on loaf volume of brown bread. *Lebensm Wiss Technol* **32**:349–356 (1999).
- 39 Ozturk S, Ozboy O, Cavidoglu I and Koksel H, Effects of brewer's spent grain on the quality and dietary fibre content of cookies. *J Inst Brew* **108**:23–27 (2002).
- 40 Uysal H, Bilgili N, Elgun A, Ibanoglu S, Herken EN and Demir MK, Effect of dietary fibre and xylanase enzyme addition on the selected properties of wire-cut cookies. *J Food Eng* **78**:1074–1078 (2007).
- 41 Kulp K, Enzymolysis of pentosans of wheat flour. *Cereal Chem* **45**:339–350 (1968).
- 42 Kim SK and D'Appolonia BL, Bread staling studies. 3. Effect of pentosans on dough, bread, and bread staling rate. *Cereal Chem* **54**:225–229 (1997).
- 43 Meuser F and Suckow P, Non-starch polysaccharides, in *Chemistry and Physics of Baking*, ed. by Blanshard JMV, Frazier PJ and Galliard T. Burlington House, London, pp. 42–61 (1986).
- 44 Fox PF and Mulvihill DM, Enzymes in wheat, flour, and bread, in *Advances in Cereal Science and Technology*, ed. by Pomeranz Y. American Association of Cereal Chemists, St Paul, MN, pp. 107–156 (1986).
- 45 Ter Haseborg E and Himmelstein A, Quality problems with high fiber breads solved with the use of hemicellulase enzymes. *Cereal Foods World* **33**:419–422 (1988).
- 46 Rouau X, El-Hayek ML and Moreau D, Effect of an enzyme preparation containing pentosanases on the bread-making quality of flours in relation to changes in pentosan properties. *J Cereal Sci* **19**:259–272 (1984).
- 47 Courtin CM, Roelants A and Delcour JA, Fractionation/reconstitution experiments provide insight into the role of endoxylanases in breadmaking. *J Agric Food Chem* **47**:1870–1877 (1999).
- 48 Courtin CM and Delcour JA, Arabinoxylans and endoxylanases in wheat flour bread-making. *J Cereal Sci* **35**:225–243 (2002).
- 49 Butt MS, Tahir-Nadeem M, Ahmad Z and Sultan MT, Xylanases and their applications in baking industry. *Food Technol Biotechnol* **46**:22–31 (2008).
- 50 Harris PJ and Smith BG, Plant cell walls and cell-wall polysaccharides: structures, properties and uses in food products. *Int J Food Sci Technol* **41**:129–143 (2006).
- 51 Inglett GE, Development of a dietary fiber gel for calorie-reduced foods. *Cereal Foods World* **42**:382–384 (1997).
- 52 Inglett GE, Dietary fiber gels for calorie reduced foods and method for preparing the same. US Patent 5766662 (1998).
- 53 Inglett GE and Carriere CJ, Cellulosic fiber gels prepared from cell walls of maize hulls. *Cereal Chem* **78**:471–475 (2001).
- 54 FNS, National school lunch program. [Online]. Available: <http://www.fns.usda.gov/cnd/Lunch/> [23 June 2009].
- 55 Warner K and Inglett GE, Flavor and texture characteristics of foods containing Z-Trim corn and oat fibers as fat or flour replacers. *Cereal Foods World* **42**:821–825 (1997).
- 56 Wolf MJ, MacMasters MM, Cannon JA, Rosewell EC and Rist CE, Preparation and some properties of hemicelluloses from corn hulls. *Cereal Chem* **30**:451–470 (1953).
- 57 Gould JM, Alkaline peroxide delignification of agricultural residues to enhance enzymatic saccharification. *Biotechnol Bioeng* **26**:46–52 (1984).
- 58 Doner LW, Chau HK, Fishman ML and Hicks KB, An improved process for isolation of corn fiber gum. *Cereal Chem* **75**:408–411 (1998).
- 59 Yadav MP, Johnston DB, Hotchkiss AT and Hicks KB, Corn fiber gum: a potential gum arabic replacer for beverage flavor emulsification. *Food Hydrocolloids* **21**:1022–1030 (2007).
- 60 Yadav MP, Fishman ML, Chau HK, Johnston DB and Hicks KB, Molecular characteristics of corn fiber gum and their influence on CFG emulsifying properties. *Cereal Chem* **84**:175–180 (2007).
- 61 Maes C and Delcour JA, Structural characterisation of water-extractable and water-unextractable arabinoxylans in wheat bran. *J Cereal Sci* **35**:315–326 (2002).
- 62 Yadav MP, Moreau RA and Hicks KB, Phenolic acids, lipids, and proteins associated with purified corn fiber arabinoxylans. *J Agric Food Chem* **55**:943–947 (2007).
- 63 Georget DMR, Ng A, Smith AC and Waldron KW, Thermal characterisation of oxidatively cross-linked American corn bran hemicellulose. *J Sci Food Agric* **79**:481–483 (1999).
- 64 Zhao ZH, Egashira Y and Sanada H, Ferulic acid is quickly absorbed from rat stomach as the free form and then conjugated mainly in liver. *J Nutr* **134**:3083–3088 (2004).
- 65 Kroon PA, Faulds CB, Ryden P, Robertson JA and Williamson G, Release of covalently bound ferulic acid from fiber in the human colon. *J Agric Food Chem* **45**:661–667 (1997).
- 66 Vitaglione P, Napolitano A and Fogliano V, Cereal dietary fibre: a natural functional ingredient to deliver phenolic compounds into the gut. *Trends Food Sci Technol* **19**:451–463 (2008).
- 67 Galvez J, Rodriguez-Cabezas ME and Zarzuelo A, Effects of dietary fiber on inflammatory bowel disease. *Mol Nutr Food Res* **49**:601–608 (2005).
- 68 King DE, Dietary fiber, inflammation, and cardiovascular disease. *Mol Nutr Food Res* **49**:594–600 (2005).
- 69 Young GP, Hu Y, Le Leu RK and Nyskohus L, Dietary fibre and colorectal cancer: a model for environment–gene interactions. *Mol Nutr Food Res* **49**:571–584 (2005).
- 70 Whistler RL, General comments on future research, in *Progress in Biotechnology I: New Approaches to Research on Cereal Carbohydrates*, ed. by Hill R and Da ML. Elsevier, New York, NY, pp. 413–415 (1985).
- 71 Vazquez MJ, Alonso JL, Dominguez H and Parajo JC, Xylooligosaccharides: manufacture and applications. *Trends Food Sci Technol* **11**:387–393 (2000).
- 72 Moure A, Gullon P, Dominguez H and Parajo JC, Advances in the manufacture, purification and applications of xylo-oligosaccharides as food additives and nutraceuticals. *Process Biochem* **41**:1913–1923 (2006).
- 73 Maes C, Vangeneugden B and Delcour JA, Relative activity of two endoxylanases towards water-unextractable arabinoxylan in wheat bran. *J Cereal Sci* **39**:181–186 (2004).
- 74 Katapodis P and Christakopoulos P, Enzymic production of feruloyl xylo-oligosaccharides from corn cobs by a family 10 xylanase from *Thermoascus aurantiacus*. *Lebensm Wiss Technol* **41**:1239–1243 (2008).
- 75 Moers K, Courtin CM, Brijs K and Delcour JA, A screening method for endo-beta-1,4-xylanase substrate selectivity. *Anal Biochem* **319**:73–77 (2003).
- 76 Berrin JG and Juge N, Factors affecting xylanase functionality in the degradation of arabinoxylans. *Biotechnol Lett* **30**:1139–1150 (2008).
- 77 Charnock SJ, Lakey JH, Virden R, Hughes N, Sinnott ML, Hazlewood GP, et al, Key residues in subsite F play a critical role in the activity of *Pseudomonas fluorescens* subspecies *Cellulosa* xylanase A against xylooligosaccharides but not against highly polymeric substrates such as xylan. *J Biol Chem* **272**:2942–2951 (1997).
- 78 Biely P, Vrsanska M, Tenkanen M and Kluepfel D, Endobeta-1,4-xylanase families: differences in catalytic properties. *J Biotechnol* **57**:151–166 (1997).

- 79 De Vries RP, Kester HCM, Poulsen CH, Benen JAE and Visser J, Synergy between enzymes from *Aspergillus* involved in the degradation of plant cell wall polysaccharides. *Carbohydr Res* **327**:401–410 (2000).
- 80 Moore J, Bamforth CW, Kroon PA, Bartolome B and Williamson G, Ferulic acid esterase catalyses the solubilization of beta-glucans and pentosans from the starchy endosperm cell walls of barley. *Biotechnol Lett* **18**:1423–1426 (1996).
- 81 Faulds CB, Mandalari G, LoCurto R, Bisignano G and Waldron KW, Arabinoxylan and mono- and dimeric ferulic acid release from brewer's grain and wheat bran by feruloyl esterases and glycosyl hydrolases from *Humicola insolens*. *Appl Microbiol Biotechnol* **64**:644–650 (2004).
- 82 Figueroa-Espinoza MC, Poulsen C, Soe JB, Zargahi MR and Rouau X, Enzymatic solubilization of arabinoxylans from native, extruded, and high-shear-treated rye bran by different endo-xylanases and other hydrolyzing enzymes. *J Agric Food Chem* **52**:4240–4249 (2004).
- 83 Vegas R, Alonso JL, Dominguez H and Parajo JC, Manufacture and refining of oligosaccharides from industrial solid wastes. *Ind Eng Chem Res* **44**:614–620 (2005).
- 84 Benko Z, Andersson A, Szegyel Z, Gaspar M, Reczey K and Stalbrand H, Heat extraction of corn fiber hemicellulose. *Appl Biochem Biotechnol* **136/140**:253–265 (2007).
- 85 Rose DJ and Inglett GE, Production of feruloylated arabinoxyloligosaccharides from maize (*Zea mays*) bran by microwave-assisted autohydrolysis. *Food Chem* **119**:1613–1618 (2010).
- 86 Rastall RA and Maitin V, Prebiotics and synbiotics: towards the next generation. *Curr Opin Biotechnol* **13**:490–496 (2002).
- 87 Gibson GR and Roberfroid MB, Dietary modulation of the human colonic microbiota – introducing the concept of prebiotics. *J Nutr* **125**:1401–1412 (1995).
- 88 Hsu CK, Liao JW, Chung YC, Hsieh CP and Chan YC, Xylooligosaccharides and fructooligosaccharides affect the intestinal microbiota and precancerous colonic lesion development in rats. *J Nutr* **134**:1523–1528 (2004).
- 89 Chung YC, Hsu CK, Ko CY and Chan YC, Dietary intake of xylooligosaccharides improves the intestinal microbiota, fecal moisture, and pH value in the elderly. *Nutr Res* **27**:756–761 (2007).
- 90 Ohta T, Yamasaki S, Egashira Y and Sanada H, Antioxidant activity of corn bran hemicellulose fragments. *J Agric Food Chem* **42**:653–656 (1994).
- 91 Yuan X, Wang J and Yao H, Antioxidant activity of feruloylated oligosaccharides from wheat bran. *Food Chem* **90**:759–764 (2005).
- 92 Wang J, Sun BG, Cao YP, Song HL and Tian Y, Inhibitory effect of wheat bran feruloyl oligosaccharides on oxidative DNA damage in human lymphocytes. *Food Chem* **109**:129–136 (2008).
- 93 Lo HH and Chung JG, The effects of plant phenolics, caffeic acid, chlorogenic acid and ferulic acid on arylamine *N*-acetyltransferase activities in human gastrointestinal microflora. *Anticancer Res* **19**:133–139 (1999).
- 94 Yuan X, Wang J and Yao H, Feruloyl oligosaccharides stimulate the growth of *Bifidobacterium bifidum*. *Anaerobe* **11**:225–229 (2005).
- 95 Zhao Z, Egashira Y and Sanada H, Digestion and absorption of ferulic acid sugar esters in rat gastrointestinal tract. *J Agric Food Chem* **51**:5534–5539 (2003).
- 96 Zhao Z and Moghadasian MH, Chemistry, natural sources, dietary intake and pharmacokinetic properties of ferulic acid: a review. *Food Chem* **109**:691–702 (2008).
- 97 White PJ and Weber EJ, Lipids of the kernel, in *Corn: Chemistry and Technology*, ed. by White PL and Johnson LA. American Association of Cereal Chemists, St Paul, MN, pp. 355–406 (2003).
- 98 Wilson TA, DeSimone AP, Romano CA and Nicolosi RJ, Corn fiber oil lowers plasma cholesterol levels and increases cholesterol excretion greater than corn oil and similar to diets containing soy sterols and soy stanols in hamsters. *J Nutr Biochem* **11**:443–449 (2000).
- 99 Ramjiganesh RST, Freake HC, McIntyre JC and Fernandez ML, Corn fiber oil lowers plasma cholesterol by altering hepatic cholesterol metabolism and up-regulating LDL receptors in guinea pigs. *J Nutr* **132**:335–340 (2002).
- 100 Jain D, Ebine N, Jia X, Kassis A, Marinangeli C, Fortin M, et al, Corn fiber oil and sitostanol decrease cholesterol absorption independently of intestinal sterol transporters in hamsters. *J Nutr Biochem* **19**:229–236 (2008).
- 101 Moreau RA, Singh V, Eckhoff SR, Powell MJ, Hicks KB and Norton RA, Comparison of yield and composition of oil extracted from corn fiber and corn bran. *Cereal Chem* **76**:449–451 (1999).
- 102 Norton R, Quantitation of steryl ferulate and *p*-coumarate esters from corn and rice. *Lipids* **30**:269–274 (1995).
- 103 Singh V, Moreau RA, Haken AE, Hicks KB and Eckhoff SR, Effect of various acids and sulfites in steep solution on yields and composition of corn fiber and corn fiber oil. *Cereal Chem* **77**:665–668 (2000).
- 104 Singh V, Johnston DB, Moreau RA, Hicks KB, Dien BS and Bothast RJ, Pretreatment of wet-milled corn fiber to improve recovery of corn fiber oil and phytosterols. *Cereal Chem* **80**:118–122 (2003).
- 105 Wu YV and Norton RA, Enrichment of protein, starch, fat, and sterol ferulates from corn fiber by fine grinding and air classification. *Ind Crop Prod* **14**:135–138 (2001).
- 106 Moreau RA, Hicks KB and Powell MJ, Effect of heat pretreatment on the yield and composition of oil extracted from corn fiber. *J Agric Food Chem* **47**:2869–2871 (1999).
- 107 Graf E, Antioxidant potential of ferulic acid. *Free Radic Biol Med* **13**:435–448 (1992).
- 108 Heinonen M, Rein D, Satue-Gracia MT, Huang SW, German JB and Frankel EN, Effect of protein on the antioxidant activity of phenolic compounds in a lecithin–liposome oxidation system. *J Agric Food Chem* **46**:917–922 (1998).
- 109 Stead D, The effect of hydroxycinnamic acids and potassium sorbate on the growth of 11 strains of spoilage yeasts. *J Appl Bacteriol* **78**:82–87 (1995).
- 110 Tsou MF, Hung CF, Lu HF, Wu LT, Chang SH, Chang HL, et al, Effects of caffeic acid, chlorogenic acid and ferulic acid on growth and arylamine *N*-acetyltransferase activity in *Shigella sonnei* (group D). *Microbios* **101**:37–46 (2000).
- 111 Ou S and Kwok KC, Ferulic acid: pharmaceutical functions, preparation and applications in foods. *J Sci Food Agric* **84**:1261–1269 (2004).
- 112 Kwok KC and Ou SY, Application of ferulic acid in the preparation of SPI-based edible film. *Sci Technol Food Ind* **129**:24–27 (2002).
- 113 O'Connell JE and Fox PF, Proposed mechanism for the effect of polyphenols on the heat stability of milk. *Int Dairy J* **9**:523–536 (1999).
- 114 Adams R and Bockstahler TE, Preparation and reaction of hydroxycinnamic acids and esters. *J Am Chem Soc* **74**:5346–5348 (1952).
- 115 Tilay A, Bule M, Kishenkumar J and Annapure U, Preparation of ferulic acid from agricultural wastes: its improved extraction and purification. *J Agric Food Chem* **56**:7644–7648 (2008).
- 116 Crepin VF, Faulds CB and Connerton IF, Functional classification of the microbial feruloyl esterases. *Appl Microbiol Biotechnol* **63**:647–652 (2004).
- 117 Wong DWS, Feruloyl esterase – a key enzyme in biomass degradation. *Appl Biochem Biotechnol* **133**:87–112 (2006).
- 118 Topakas E, Vafiadi C and Christakopoulos P, Microbial production, characterization and applications of feruloyl esterases. *Process Biochem* **42**:497–509 (2007).
- 119 Mathew S and Abraham TE, Studies on the production of feruloyl esterase from cereal brans and sugar cane bagasse by microbial fermentation. *Enzym Microb Technol* **36**:565–570 (2005).
- 120 Faulds CB, Kroon PA, Saulnier L, Thibault JF and Williamson G, Release of ferulic acid from maize bran and derived oligosaccharides by *Aspergillus niger* esterases. *Carbohydr Polym* **27**:187–190 (1995).
- 121 Saulnier L, Marot C, Elgorriaga M, Bonnin E and Thibault JF, Thermal and enzymatic treatments for the release of free ferulic acid from maize bran. *Carbohydr Polym* **45**:269–275 (2001).
- 122 Ralet MC, Faulds CB, Williamson G and Thibault JF, Degradation of feruloylated oligosaccharides from sugar-beet pulp and wheat bran by ferulic acid esterases from *Aspergillus niger*. *Carbohydr Res* **263**:257–269 (1994).
- 123 Aminoff C, Vanninen E and Doty TE, The occurrence, manufacture and properties of xylitol, in *Xylitol*, ed. by Counsell JN. Applied Science Publishers, London, pp. 1–9 (1978).
- 124 Winklenlausen E and Kuzmanova S, Microbial conversion of D-xylose to xylitol. *J Ferment Bioeng* **86**:1–14 (1998).
- 125 Wenzl HFJ, *The Chemical Technology of Wood*. Academic Press, New York, NY, pp. 92–156 (1970).

- 126 Rao RS, Jyothi CP, Prakasham RS, Sarma PN and Rao LV, Xylitol production from corn fiber and sugarcane bagasse hydrolysates by *Candida tropicalis*. *Bioresource Technol* **97**:1974–1978 (2006).
- 127 Buhner J and Agblevor FA, Effect of detoxification of dilute-acid corn fiber hydrolysate on xylitol production. *Appl Biochem Biotechnol* **119**:13–30 (2004).
- 128 Esposito LJ, Formanek K, Kientz G, Mauger F, Maureaux V, Robert G, *et al*, Vanillin, in *Encyclopedia of Chemical Technology* (4th edn), Vol. 24, ed. by Kroschwitz JI and Howe-Grant M. Wiley, New York, NY, pp. 813–825 (1997).
- 129 Hocking MB, Vanillin: synthetic flavoring from spent sulfite liquor. *J Chem Educ* **74**:1055–1059 (1997).
- 130 Goodner KL, Jella P and Rouseff RL, Determination of vanillin in orange, grapefruit, tangerine, lemon, and lime juices using GC–olfactometry and GC–MS/MS. *J Agric Food Chem* **48**:2882–2886 (2000).
- 131 Buranov AU and Mazza G, Extraction and purification of ferulic acid from flax shives, wheat and corn bran by alkaline hydrolysis and pressurised solvents. *Food Chem* **115**:1542–1548 (2009).
- 132 Dignum MJW, Kerler J and Verpoorte R, Vanilla production: technological, chemical, and biosynthetic aspects. *Food Rev Int* **17**:199–219 (2001).
- 133 Rosazza JPN, Huang Z, Dostal L, Volm T and Rousseau B, Biocatalytic transformations of ferulic acid: an abundant aromatic natural product. *J Ind Microbiol* **15**:457–471 (1995).
- 134 Walton NJ, Narbad A, Faulds CB and Williamson G, Novel approaches to the biosynthesis of vanillin. *Curr Opin Biotechnol* **11**:490–496 (2000).
- 135 Mathew S and Abraham TE, Bioconversions of ferulic acid, an hydroxycinnamic acid. *Crit Rev Microbiol* **32**:115–125 (2006).
- 136 Priefert H, Rabenhorst A and Steinbuchel A, Biotechnological production of vanillin. *Appl Microbiol Biotechnol* **56**:296–314 (2001).
- 137 Lesage-Meessen L, Lomascolo A, Bonnin E, Thibault JF, Buleon A, Roller M, *et al*, A biotechnological process involving filamentous fungi to produce natural crystalline vanillin from maize bran. *Appl Biochem Biotechnol* **102**:141–153 (2002).